Does prematurity moderate the heritability of early motor development?

Catharina E. M. van Beijsterveldt, Cyrina B. Brouwer, Meike Bartels & Dorret I. Boomsma Netherlands Twin Registry (NTR) VU Amsterdam

Introduction: Motor milestones are recognizable landmarks of development in the first years of life and good indicators of early motor development. One risk factor for a delayed motor development is prematurity. In this study we estimated the contribution of genetic factors on achievement of motor milestones and tested whether prematurity moderates these factors. Because of immaturity of organs and systems in preterm twins we expect that genetic factors contribute to a greater extent in the variation of achieving motor milestones in term born twins than in preterm born twins.

mijlpalen

The study is part of the longitudinal data collection of the Young-Netherlands Twin Registry on behavioral and emotional development. Data on motor development and gestational age were available for 21,366 twin pairs, born between 1987 and 2010, and obtained by maternal report (see right). Motor development was assessed by the age of achieving a motor milestone (turning over, sitting up, crawling, standing, and walking).

With structural equation model, we estimated the effects of genetic and environmental influences and tested the moderation effect of prematurity and sex on the means and the variance components.

Examples of the zitten zonder steun? questions on motor milestones kruipen op handen en knieën?

Distribution of prematurity

Very preterm (<32 wks)	926 pairs (4.3%)
Preterm (>=32 wks and <37 wks)	7520 pairs (35.2%)
Term (>= 37 wks)	12920 pairs (60.5%)

Genetic and environmental influences

Standardized estimates

Unstandardized estimates

turning

Results

Mean and SD for motor milestones

turning





Conclusion

- Effect of prematurity was highly significant for all motor milestones.
- Except for turning, boys achieved motor milestones later than girls. However effect sizes were small, varying from 2 to 5 days.
- Individual differences in motor milestones are explained by both genetic and shared environmental influences.
- Heritability estimates are highest for walking and lowest for sitting. • Standardized heritability estimates were higher in term born twins than in (very) preterm born. Inspection of the unstandardized estimates showed that the higher heritability could be explained by an increase of the shared environmental variance in the (very) preterm born twins. Genetic variance did not change across groups.

