

Two-locus Linkage Analysis Applied to Putative Quantitative Trait Loci for Lipoprotein(a) Levels

Marian Beekman¹, Bastiaan T. Heijmans¹, Nicholas G. Martin², John B. Whitfield³, Nancy L. Pedersen⁴, Ulf DeFaire⁵, Harold Snieder^{6,7}, Nico Lakenberg¹, Peter de Knijff⁸, Rune R. Frants⁸, Gert Jan B. van Ommen⁸, Cornelis Klufft⁹, George P. Vogler¹⁰, P. Eline Slagboom¹, and Dorret I. Boomsma¹¹

¹ Section of Molecular Epidemiology, Leiden University Medical Center, Leiden, The Netherlands

² Queensland Institute for Medical Research, Brisbane, Australia

³ Department of Clinical Biochemistry, Royal Prince Alfred Hospital, Sydney, Australia

⁴ Department of Medical Epidemiology, Karolinska Institute, Stockholm, Sweden

⁵ Institute of Environmental Medicine, Karolinska Institute, Stockholm, Sweden

⁶ Twin Research & Genetic Epidemiology Unit, St. Thomas' Hospital, London, United Kingdom

⁷ Georgia Prevention Institute, Department of Pediatrics, Medical College of Georgia, Augusta, United States of America

⁸ Department of Human and Clinical Genetics, Leiden University Medical Center, Leiden, The Netherlands

⁹ Department of Vascular and Connective Tissue Research, TNO Prevention and Health, Leiden, The Netherlands

¹⁰ Department of Biobehavioral Health, Center for Developmental and Health Genetics, Pennsylvania State University, University Park, Pennsylvania, United States of America

¹¹ Department of Biological Psychology, Free University of Amsterdam, Amsterdam, The Netherlands

Plasma levels of lipoprotein(a) — Lp(a) — are associated with cardiovascular risk (Danesh et al., 2000) and were long believed to be influenced by the *LPA* locus on chromosome 6q27 only. However, a recent report of Broeckel et al. (2002) suggested the presence of a second quantitative trait locus on chromosome 1 influencing Lp(a) levels. Using a two-locus model, we found no evidence for an additional Lp(a) locus on chromosome 1 in a linkage study among 483 dizygotic twin pairs.

Levels of Lp(a) are highly heritable. More than 90% of the variance is attributable to genetic factors (Rainwater, 1996; Snieder et al., 1997). Variation in the gene encoding apolipoprotein(a) (*LPA*) on chromosome 6q27 accounts for more than 85% of the variation in Lp(a) plasma levels (Boerwinkle et al., 1992; Boomsma et al., 1993; DeMeester et al., 1995). Recently, Broeckel et al. (2002) reported a second chromosomal region linked with Lp(a) levels on chromosome 1 at 170 cM from pter, which explained 16% of the variance in Lp(a), in a linkage study of 513 Western European families with 2 or more cases of premature myocardial infarction (on average 2.7 subjects per family). To test whether this putative quantitative trait locus (QTL) also affects Lp(a) levels in the general population, we performed a linkage study in four samples of dizygotic (DZ) and monozygotic (MZ) twin pairs recruited regardless of their health status.

Materials and Methods

Plasma levels of Lp(a) were obtained of an adolescent Dutch sample (90 DZ, 70 MZ, mean age = 17 years) and three adult samples from the Netherlands (119 DZ, 94 MZ, mean age = 44 years), Sweden (87 DZ, 69 MZ, mean

age = 65 years) and Australia (669 DZ, 734 MZ, mean age = 44 years) recruited regardless of their health status (Beekman et al., 2002). In these samples, heritability estimates of Lp(a) levels ranged from 90 to 96%. Genotypes for the linkage study were obtained for DZ pairs for whom DNA was available, and not for MZ twins, since these are genetically identical, sharing two alleles identical-by-descent (IBD) at all loci. Markers from the Marshfield Screening Set 8 were genotyped on chromosome 1 and 6 with an average spacing of 18 cM (Beekman et al., 2001). To adjust for non-normality, Lp(a) levels were transformed by the natural logarithm. Variance components linkage analyses were carried out, including data from the 483 genotyped DZ twins as well as the phenotypical data from the 967 untyped MZ twin pairs. The inclusion of MZ twins provides a more accurate estimate of non-shared environmental effects (MZ correlations are < 1), so that an upper limit for the estimation of genetic effects is obtained and overestimation is reduced. IBD status for the DZ pairs was estimated separately for each of the four samples with Genehunter 2.1 (Kruglyak et al., 1996) using population specific allele frequencies. All analyses were performed using Mx 1.52d (Neale et al., 1999) in a 4-sample simultaneous analysis. Mean Lp(a) levels, background genetic and non-shared environmental effects were estimated for each sample separately. The total variance was modeled as $A + E + Q$, where *A* represents additive genetic background

Address for correspondence: Marian Beekman, Section of Molecular Epidemiology, Leiden University Medical Center, PO Box 9503, 2300 RA, Leiden, The Netherlands. Email: M.Beekman@lumc.nl

factors, E non-shared environmental factors and Q the QTL effect. The covariance equals $A + Q$ for MZ twins and $\frac{1}{2}A + \hat{\pi}Q$ for DZ twins, where $\hat{\pi}$ is estimated as $\frac{1}{2}P(\text{IBD} = 1) + P(\text{IBD} = 2)$. To test for linkage, the fit of the AE model was compared to the fit of the AEQ model. For the two-locus models, the equations for the total variance, covariance for MZ twins and covariance for DZ twins were $A + E + Q_1 + Q_6$, $A + Q_1 + Q_6$ and $\frac{1}{2}A + \hat{\pi}_1Q_1 + \hat{\pi}_6Q_6$, respectively, where Q_1 and Q_6 refer to the putative QTLs on chromosomes 1 and 6, respectively. The proportion of the variance explained by the QTL was constrained to be equal over the samples. The power to replicate the putative Lp(a) QTL on chromosome 1 (QTL effect, 0.16; residual shared variance, 0.74) in 483 sib pairs at a significance level of .00074 (LOD-score of 2.2 constituting suggestive linkage according to Lander-Kruglyak criteria) and assuming incomplete marker informativeness ($\theta = 0.10$) is 0.51, as calculated using the Genetic Power Calculator (<http://statgen.iop.kcl.ac.uk/gpc/>) (Sham et al., 2000).

Results

As expected, we obtained highly significant evidence for linkage at the *LPA* locus on chromosome 6 (maximum LOD score {MLS} = 9.8, Figure 1; MLS of 1.7, 4.4, 1.8 and 5.1, in the adolescent Dutch and adult Dutch, Swedish and Australian samples, respectively). This QTL explained 82% of the total variance in Lp(a) levels (Table 1). Next, chromosome 1 was analyzed and a MLS of 1.6 was

obtained at 251 cM from pter (Figure 1; MLS of 0.0, 0.7, 2.2 and 1.8, in the adolescent Dutch and adult Dutch, Swedish and Australian samples, respectively).

This QTL explained 44% of the total variance in Lp(a) levels (Table 1). The two putative QTLs on chromosome 6 and 1 together explained 126% of the variation in Lp(a) levels, which likely results from overestimation of effect sizes, a common phenomenon in genome-wide linkage analyses (Göring et al., 2001). We, therefore, analyzed the effect of both QTLs simultaneously in a two-locus model (Figure 1). In this two-locus model, evidence for linkage on chromosome 1 completely disappeared (LOD = 0; Figure 1). Moreover, the chromosome 1 locus now explained none of the variation in Lp(a) plasma levels, while the estimates for the *LPA* locus remained unaffected (Table 1).

Conclusion

Although our first analysis indicated linkage with Lp(a) on chromosome 1, disappearance of this linkage in a two-locus model suggests that, besides a possible power problem to detect an additional QTL with a small effect, our initial finding may have been a false positive result. Our analyses thus indicate that multi-locus models as employed here, may be a useful tool to distinguish true linkage results from spurious ones. In a recent study, Broeckel et al. (2002) reported a second QTL on chromosome 1 influencing Lp(a) levels in premature MI patients. Our study in twin pair samples provides no support for a Lp(a) QTL on chromosome 1 acting in the general population.




Figure 1

Linkage analysis of chromosome 6 and 1 with plasma levels of Lp(a) in four twin samples simultaneously; — shows the result of the initial analysis of the chromosomes; --- shows the result of the two-locus analysis.

Table 1

Effect Size of Putative Lp(a) QTLs at Chromosome 6 and 1 at the Position of the Maximum LOD Scores, in Separate and Simultaneous Analyses

Chromosome analyzed	Maximum LOD score (position in cM from pter)		Proportion of the variance attributable to the QTL at chromosome 6 (95% C.I.)	Proportion of the variance attributable to the QTL at chromosome 1 (95% C.I.)
	Chr 6	Chr 1		
6	9.8 (163)	—	0.82 (0.66–0.91)	—
1	—	1.6 (251)	—	0.44 (0.13–0.68)
6 and 1	0.82 (163)	0.0 (251)	0.82 (0.65–0.91)	0.00 (0.00–0.14)

Acknowledgments

This study was supported by grants to G.P. Vogler of the National Institutes of Health, National Heart, Lung, and Blood Institute (HL55976), to N.G. Martin and J.B. Whitfield of the National Heart Foundation of Australia, to N.L. Pedersen (AG04563) of NIA, to PE Slagboom (98.199) and D.I. Boomsma (86.083, 88.042 and 90.313) of the Netherlands Heart Foundation.

References

- Beekman, M., Heijmans, B. T., Martin, N. G., Pedersen, N. L., Whitfield, J. B., DeFaire, U., et al. (2002). Heritabilities of apolipoprotein and lipid levels in three countries. *Twin Research*, *5*(2), 87–97.
- Beekman, M., Lakenberg, N., Cherny, S. S., de Knijff, P., Kluff, C. C., van Ommen, G. J., et al. (2001). A powerful and rapid approach to human genome scanning using small quantities of genomic DNA. *Genetical Research*, *77*(2), 129–134.
- Boerwinkle, E., Leffert, C. C., Lin, J., Lackner, C., Chiesa, G., & Hobbs, H. H. (1992). Apolipoprotein(a) gene accounts for greater than 90% of the variation in plasma lipoprotein(a) concentrations. *Journal of Clinical Investigation*, *90*(1), 52–60.
- Boomsma, D. I., Kaptein, A., Kempen, H. J., Gevers, L. J., & Princen, H. M. (1993). Lipoprotein(a): Relation to other risk factors and genetic heritability. Results from a Dutch parent-twin study. *Atherosclerosis*, *99*(1), 23–33.
- Brockel, U., Hengstenberg, C., Mayer, B., Holmer, S., Martin, L. J., Comuzzie, A. G., et al. (2002). A comprehensive linkage analysis for myocardial infarction and its related risk factors. *Nature Genetics*, *30*(2), 210–214.
- Danesh, J., Collins, R., & Peto, R. (2000). Lipoprotein(a) and coronary heart disease. Meta-analysis of prospective studies. *Circulation*, *102*(10), 1082–1085.
- DeMeester, C. A., Bu, X., Gray, R. J., Lusia, A. J., & Rotter, J. I. (1995). Genetic variation in lipoprotein (a) levels in families enriched for coronary artery disease is determined almost entirely by the apolipoprotein (a) gene locus. *American Journal of Human Genetics*, *56*(1), 287–293.
- Göring, H. H., Terwilliger, J. D., & Blangero, J. (2001). Large upward bias in estimation of locus-specific effects from genomewide scans. *American Journal of Human Genetics*, *69*(6), 1357–1369.
- Kruglyak, L., Daly, M. J., Reeve-Daly, M. P., & Lander, E. S. (1996). Parametric and nonparametric linkage analysis: A unified multipoint approach. *American Journal of Human Genetics*, *58*(6), 1347–1363.
- Neale, M. C., Boker, S. M., Xie, G., & Maes, H. H. (1999). *Mx: Statistical modeling* (5th ed.). Richmond, VA: VCU, Department of Psychiatry.
- Rainwater, D. L. (1996). Lp(a) concentrations are related to plasma lipid concentrations. *Atherosclerosis*, *127*(1), 13–18.
- Sham, P. C., Cherny, S. S., Purcell, S., & Hewitt, J. K. (2000). Power of linkage versus association analysis of quantitative traits, by use of variance-components models, for sibship data. *American Journal of Human Genetics*, *66*(5), 1616–1630.
- Snieder, H., van Doornen, L. J., & Boomsma, D. I. (1997). The age dependency of gene expression for plasma lipids, lipoproteins, and apolipoproteins. *American Journal of Human Genetics*, *60*(3), 638–650.